

Measuring Elevation Change in Meadowlands Marshes Using Surface Elevation Tables (SETs) and Marker Horizons

Meadowlands Research and Restoration Institute (December, 2022)

Introduction

Sediment Elevation Tables (SET) provide a constant plane in space from which the change in marsh surface elevation can be measured. Benchmark rods and marker horizons of feldspar were established at five sites in the lower Hackensack River Meadowlands during August of 2008. The five sites being monitored are Riverbend High Marsh (RBH), Riverbend Mixed Marsh (RBM), Sawmill (SM), Lyndhurst Riverside (LR), and Secaucus High School (SHS). Each site is measured annually, except in the year 2009. This Report is a summary of the marsh elevation change up to today. (Fall 2022)

Figure 1: Location of the SET sites




Methods:

Three replicate plots are installed at each of the five sites. At each plot within a site, a total of 36 measurements are made covering the four cardinal directions, which gives a total of 108 measurements per site. Every year, measurements in each site are compared to the previous year and this constitutes a data point. To obtain the rate of elevation change, the last data point value is divided by the number of years elapsed between the establishment of the site. Time elapsed between readings is summarized in this report (Table 1).

Diagram of the SET benchmark plot:

The four outside corners, A, B, C, and D represent feldspar horizon markers. The yellow circle inside with a, b, c, and d are the four cardinal directions in which the measurements are taken.



Locations were chosen to span several miles of tidal wetlands and represent different vegetation and marsh regimes. The five sites selected include a restored *Spartina alterniflora* low marsh (Secaucus High School, SHS), a *Spartina alterniflora* low marsh (Saw Mill, SM), a *Spartina patens* dominated high marsh (Riverbend-Patens, RBP), a mixed *Spartina patens* and *Phragmites australis* high marsh (Riverbend Mixed, RBM) and a *Phragmites australis* dominated high marsh (Lyndhurst Riverside, LR). At each site, three replicate plots were installed. At each plot, nine pins are lowered to the marsh surface. Readings are taken in each of four directions resulting in a total of 108 measurements for each of the 5 sites. At the time of each subsequent reading, results obtained from each pin are compared. The average of the resulting differences becomes one data point that represents the level of the marsh surface elevation.

Table 1: Time Elapsed Between SET installation the latest sampling

Location	Initial Date	Most Recent Date	Days	Years
Riverbend High/Mixed Marsh	8/26/2008	11/21/2022	5200	14.247
Sawmill	8/28/2008	12/1/2022	5208	14.27
Lyndhurst Riverside	8/29/2008	12/1/2022	5207	14.27
Secaucus High School	8/28/2008	11/22/2022	5199	14.24

Table 2 Averages of elevation changes obtained at each of the five sites. The average of all the plots is then divided by the time elapsed since the initial date (Table 1) to obtain the rate of elevation change in mm/yr (Table 2a).

Riverbend High Marsh (RBH)		Riverbend Mixed Marsh (RBM)		Sawmill		Lyndhurst (LR)		Secaucus HS (SHS)	
All Plots (mm)	72.01	All Plots (mm)	69.34	All Plots (mm)	91.72	All Plots (mm)	52.32	All Plots (mm)	86.12
Std Error	4.49	Std Error	11.83	Std Error	2.43	Std Error	4.12	Std Error	3.42
Plot 1 (mm)	74.40	Plot 1 (mm)	48.15	SM-1 (mm)	96.47	SM-1 (mm)	44.89	SM-1 (mm)	92.3
Std Error	0.99	Std Error	7.76	Std Error	11.36	Std Error	9.65	Std Error	20.99
Elevation rate mm/yr		Elevation rate mm/yr		Elevation rate (mm/yr)		Elevation rate mm/yr		Elevation rate mm/yr	
5.05		4.87		6.93		3.95		6.05	

Table 2a: SETs Elevation Rates, Locations and Marsh type - Fall 2022 sampling

Location	Marsh Type	Dominant Vegetation	Rate of Elevation Change from 2008 to 2022 (mm/yr.)
Riverbend Patens (RBP)	High Marsh	<i>Spartina patens</i>	5.1
Riverbend Mixed (RBM)	High Marsh	<i>Phragmites australis</i> / <i>Spartina patens</i>	4.9
Sawmill (SM)	High Marsh	<i>Spartina alterniflora</i>	6.4
Lyndhurst Riverside (LR)	High Marsh	<i>Phragmites australis</i>	4.0
Secaucus (SHS)	Low Marsh	<i>Spartina alterniflora</i>	6.1

Feldspar horizons were placed inside three corners of each plot. The sediment between the white feldspar marker and the horizon is measured. One reading is taken at each of the three corners resulting in a total of nine values per site. The average of all readings is shown in Table 3. All recoverable values are included in the calculation for accretion rate. To obtain a yearly rate, total accretion is divided by the number of years that have elapsed between establishment of the benchmark and the latest reading. Fourteen years elapsed between the readings summarized in this report. Table 1 provides the dates for each reading and the time elapsed in days and years.

Table 3: Average Accretion Tables (mm) - Fall 2022 sampling

Riverbend High Marsh (RBH)		Riverbend Mixed Marsh (RBM)		Sawmill		Lyndhurst Riverside (LR)		Secaucus HS (SHS)	
All Plots (mm)	63.33	All Plots (mm)	81.67	All Plots (mm)	58.33	All Plots (mm)	53.33	All Plots (mm)	78.3
Std Error	6.67	Std Error	7.26	Std Error	1.67	Std Error	4.41	Std Error	1.88
SM-1 (mm)	70.0	RBM-1 (mm)	95.00	TM-1 (mm)	60.00	SM-1 (mm)	45.00	SM-1 (mm)	110.0
Std Error	0.00	Std Error	0.00	Std Error	0.00	Std Error	0.00	Std Error	0.00
SM-2 (mm)	50.0	RBM-2 (mm)	80.00	TM-2 (mm)	60.00	SM-2 (mm)	60.00	SM-2 (mm)	80.0
Std Error	0.00	Std Error	0.00	Std Error	0.00	Std Error	0.00	Std Error	0.00
SM-3 (mm)	70.00	RBM-3 (mm)	70.00	TM-3 (mm)	55.00	SM-3 (mm)	55.00	SM-3 (mm)	45.0
Std Error	0.00	Std Error	0.00	Std Error	0.000	Std Error	0.00	Std Error	0.00

Table 3a: Average Accretion Rate - Fall 2022 sampling

Location	Accretion rate (mm/yr)
Riverbend High Marsh (RBH)	4.5
Riverbend Mixed Marsh (RBM)	5.7
Sawmill (SM)	7.6
Lyndhurst Accretion (LR)	3.7
Secaucus HS (SHS)	5.5

**Table 4: Summary of Elevation and Accretion rate up to the Fall of 2022.
Eight Day Swamp, Walden Swamp, Tollgate Marsh, Lyndhurst Riverside and Secaucus High School**

Riverbend High Marsh (RBH)													
Days	0	632	980	1555	2065	2430	2795	3137	3585	4045	4468	4811	5200
Sample Date	8/26/08	5/20/10	5/3/11	11/28/12	4/22/14	4/22/15	4/21/16	3/29/17	6/20/18	9/23/19	11/19/20	10/28/21	11/21/22
Elevation Rate mm/yr	0.00	2.48	6.02	5.92	3.68	4.18	3.02	4.18	4.00	5.00	3.69	5.04	5.05
Accretion Rate mm/yr	0.00	0.00	5.69	6.39	5.03	5.67	6.31	3.81	5.75	5.36	5.04	6.41	4.45

Riverbend Mixed Marsh (RBM)													
Days	0	632	980	1555	2065	2430	2795	3137	3585	4045	4468	4811	5200
Sample Date	8/26/08	5/20/10	5/3/11	11/28/12	4/22/14	4/22/15	4/21/16	3/29/17	6/20/18	9/23/19	11/19/20	10/28/21	11/21/22
Elevation Rate mm/yr	0.00	5.96	7.58	6.99	3.63	5.15	2.94	4.08	3.67	4.10	3.86	4.52	4.87
Accretion Rate mm/yr	0.00	0.00	7.80	6.31	5.05	7.18	8.59	3.75	5.88	5.21	4.81	6.15	5.73

Sawmill (SM)													
Days	0	631	980	1566	2065	2430	2814	3136	3584	4047	4467	4828	5208
Sample Date	8/28/08	5/21/10	5/5/11	12/11/12	4/24/14	4/24/15	5/12/16	3/30/17	6/21/18	9/27/19	11/20/20	11/16/21	12/1/22
Elevation Rate mm/yr	0.00	-4.66	5.51	7.94	5.70	4.74	3.93	4.75	3.94	4.20	4.65	5.06	6.43
Accretion Rate mm/yr	0.00	0.00	13.80	11.10	9.90	9.70	7.40	8.50	8.50	7.80	-	6.55	7.55

Lyndhurst Riverside (LR)													
Days	0	630	979	1557	2064	2429	3135	3583	4044	4466	4829	5207	
Sample Date	8/29/08	5/21/10	5/5/11	12/3/12	4/24/14	4/24/15	3/30/17	6/21/18	9/25/19	11/20/20	11/18/21	12/1/22	
Elevation Rate mm/yr	0	8.79	6.38	7.53	3.65	3.18	1.85	2.44	3.03	2.48	3.95	4.05	
Accretion Rate mm/yr	0	2.74	3.94	4.20	3.40	4.40	4.50	3.26	3.61	3.00	3.07	3.74	

Secaucus High School (SHS)													
Days	0	609	984	1553	2066	2421	2801	3169	3585	4044	4470	4843	5199
Sample Date	8/28/08	4/29/10	5/9/11	11/28/12	4/25/14	4/15/15	4/29/16	5/2/17	6/22/18	9/24/19	11/23/20	12/1/21	11/22/22
Elevation Rate mm/yr	0.00	9.97	5.03	6.52	6.16	5.84	3.67	3.90	3.93	3.55	9.32	10.37	6.05
Accretion Rate mm/yr	0.00	10.00	4.40	6.70	8.90	7.90	5.90	6.90	5.83	5.52	4.99	5.11	5.50

Conclusion

In the Fall of 2022, Riverbend High Marsh, or Riverbend Patens, showed an annual elevation change of 5.1 mm/yr and an accretion rate of 4.5 mm/yr. Riverbend Mixed Marsh showed an annual elevation change of 4.9 mm/yr and an accretion rate of 5.7 mm/yr. Sawmill showed an annual elevation change of 6.4 mm/yr and an accretion rate of 7.6 mm/yr. Lyndhurst Riverside, showed an annual elevation change of 4.1 mm/yr and an accretion rate of 3.7 mm/yr. Secaucus High School showed an annual elevation change of 6.1 mm/yr and an accretion rate of 5.5 mm/yr.